

Carbonate-silicate ratio for soil correction and influence on nutrition, biomass production and quality of palisade grass

Renato Ferreira de Souza^{1*}, Fabrício William Ávila², Valdemar Faquin², Adélia Aziz Alexandre Pozza², Janice Guedes Carvalho², Antônio Ricardo Evangelista³

¹EPAMIG/URECO, C.P. 295 - 35701-970 - Prudente de Morais, MG - Brasil.

²UFLA - Depto. de Ciência do Solo, C.P. 3037 - 37200-000 - Lavras, MG - Brasil.

³UFLA - Depto. de Zootecnia.

*Corresponding author <souzarf@epamig.br>

Edited by: Luís Reynaldo Ferraciu Alleoni

ABSTRACT: Silicates can be used as soil correctives, with the advantage of being a source of silicon, a beneficial element to the grasses. However, high concentrations of silicon in the plant would affect the digestibility of the forage. To evaluate the influence of the substitution of the calcium carbonate by calcium silicate on the nutrition, biomass production and the feed quality of the palisade grass [*Urochloa brizantha* (C. Hochstetter ex A. Rich.) R. Webster], three greenhouse experiments were conducted in completely randomized designs with four replications. Experimental units (pots) contained a clayey dystrophic Rhodic Haplustox, a sandy clay loam dystrophic Typic Haplustox and a sandy loam dystrophic Typic Haplustox. Each soil received substitution proportions (0, 25, 50, 75 and 100 %) of the carbonate by calcium silicate. The increase in the proportion of calcium silicate elevated the concentrations and accumulations of Si, Ca, Mg, and B, reduced Zn and did not alter P in the shoot of plants. The effects of the treatments on the other nutrients were influenced by the soil type. Inclusion of calcium silicate also increased the relative nutritional value and the digestibility and ingestion of the forage, while the concentration and accumulation of crude protein and the neutral detergent and acid detergent fibers decreased. Biomass production and feed quality of the palisade grass were generally higher with the 50 % calcium silicate treatment.

Keywords: *Urochloa brizantha*, pasture, lime, silicon, feed value

Introduction

The Brazilian savannah (cerrado) includes approximately 200 million hectares, 25 % of which are pastures, the majority being *Urochloa* spp. Between 70 to 80 % of these pastures are at some level of degradation and low productivity (Reis et al., 2010) due to soil limitations such as aluminum toxicity, phosphorus deficiency, and low pH.

Lime application has been the main method for acidity correction because of its low cost, ease of application and increase of plant availability of Ca and Mg. However, another alternative method used for the acidity correction is the application of silicates, silicon (Si) being also a source for several crops.

Authors have shown beneficial effects of Si in several vegetable species, especially grasses. Benefits include increased availability of P and micronutrients in the soil (Marschner, 1995), resistance to pests and diseases (Rodrigues et al., 2004) and tolerance to excess of Fe²⁺ and Al³⁺ ions (Mengel and Kirkby, 2001), tolerance to salinity (Matoh et al., 1986) and a more erect disposition of leaves, which increases the photosynthetic rate (Ávila et al., 2010; Yoshida et al., 1962). However, the excess uptake of Si by grasses may provide a higher mechanical resistance to degradation (Melo et al., 2003); reducing their ruminal digestibility (Jones and Handreck, 1967).

For these reasons, the objective of this study was to evaluate the effect of the substitution of calcium carbon-

ate by calcium silicate on the nutrition, biomass production and feed quality of palisade grass cultivated in soils with different texture and mineralogical composition.

Materials and Methods

Three greenhouse experiments were conducted in Lavras, state of Minas Gerais - Brazil, with three types of soil, each comprising a specific experiment. In order to include a wide variation soils with varying physical, chemical and mineralogical attributes were used: sandy loam dystrophic Typic Haplustox (LVAd-1), a sandy clay loam dystrophic Typic Haplustox (LVAd-2), and clayey dystrophic Rhodic Haplustox (LVd) (Embrapa, 2006). They were collected from the 0-20 cm layer, under natural vegetation, after the removal of the organic remains present on the surface. Later, after sieving through 4-mm mesh sieve, soil subsamples were characterized chemically, physically and mineralogically (Table 1).

Treatments were distributed in a completely randomized design with four replications and included five substitution levels (0, 25, 50, 75 and 100 %) of CaCO₃ and MgCO₃ by CaSiO₃ and MgSiO₃, and an untreated control. The salts (pa grade) were expressed in CaCO₃ equivalent (Alcarde and Rodella, 1996). In all of the treatments a 4:1 stoichiometric ratio of Ca:Mg was maintained and supplied in amounts sufficient to elevate the base saturation of each soil to 60 % (Van Raij et al., 1996) (Table 2).

After the application of the treatments and 30 days of incubation, each experimental unit received application of

Table 1 – Chemical, physical and mineralogical attributes of the soil samples (0-20 cm depth), prior to treatments.

Soil	Chemical ⁽¹⁾															
	pH	P	K	Zn	Cu	Mn	Fe	EP	Ca	Mg	Al	H+Al	T	m	V	MPAC
	mg dm ⁻³					mg L ⁻¹			mmol _c dm ⁻³ of soil				%		mg kg ⁻¹	
LVAd-1	5.2	2.0	74	2.4	0.9	7.3	86	24.4	5	2	6	40	49	40	18.4	396
LVAd-2	4.9	2.3	52	5.5	1.3	8.2	176	17.4	8	2	8	63	74	41	15.3	776
LVd	4.7	1.4	31	2.0	1.8	6.8	95	8.3	4	1	0	110	116	62	5.3	1275
	Physical ⁽²⁾															
	Sand				Silt				Clay				OM			
	g kg ⁻¹															
LVAd-1	690				80				230				19			
LVAd-2	520				80				400				26			
LVd	250				60				690				41			
	Mineralogical ⁽³⁾															
	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	TiO ₂	P ₂ O ₅	Fe _d	Fe _o	Ct	Gb	Ki	Kr					
	g kg ⁻¹ of clay															
LVAd-1	95.1	97.4	36.2	6.2	0.0	10.8	0.1	752.0	63.0	0.98	0.71					
LVAd-2	137.6	203.9	53.9	11.2	0.4	44.5	5.8	535.6	202.3	0.67	0.53					
LVd	129.8	319.1	171.8	22.0	0.9	101.6	9.2	292.7	358.8	0.41	0.26					

⁽¹⁾pH in water, Ca, Mg, K and Al were determined according to Embrapa (1997); P in the equilibrium solution (EP) according to Alvarez V. et al. (2000); H+Al and level of organic matter (OM) according to Van Raij et al. (1987). T = Cation exchange capacity at pH 7.0; m = Aluminum saturation index; V = Base saturation index and MPAC = maximum P adsorption capacity. ⁽²⁾The soil granulometry was determined by the pipette method (Day, 1965). ⁽³⁾SiO₂, Al₂O₃, Fe₂O₃, TiO₂ and P₂O₅ were determined according to Embrapa (1997); Fe_d, according to Mehra and Jackson (1960); Fe_o, according to Schwertmann (1964) and Ct (kaolinite) and Gb (gibbsite) according to Klug and Alexander (1974). Ki = SiO₂ / Al₂O₃ and Kr = SiO₂ / (Al₂O₃ + Fe₂O₃).

Table 2 – Corrective levels to elevate the base saturation of each soil to 60 %, expressed as CaCO₃ equivalent: CaCO₃ = 1.000; MgCO₃ = 1.183; CaSiO₃ = 0.879 and MgSiO₃ = 1.017.

Soil	Treatment	Corrective				Ca	Mg	Si
		CaCO ₃	MgCO ₃	CaSiO ₃	MgSiO ₃			
		%CaSiO ₃						
LVAd-1	0	815	172	0	0	320	49	0
LVAd-1	25	612	129	232	50	320	49	70
LVAd-1	50	408	86	464	100	320	49	140
LVAd-1	75	204	43	696	150	320	49	210
LVAd-1	100	0	0	928	200	320	49	280
LVAd-2	0	1323	280	0	0	519	79	0
LVAd-2	25	992	210	376	81	519	79	114
LVAd-2	50	662	140	753	163	519	79	227
LVAd-2	75	331	70	1129	244	519	79	341
LVAd-2	100	0	0	1505	325	519	79	455
LVd	0	2538	536	0	0	996	151	0
LVd	25	1904	402	722	156	996	151	218
LVd	50	1269	268	1444	312	996	151	436
LVd	75	635	134	2166	468	996	151	654
LVd	100	0	0	2887	624	996	151	873

fertilizers as basal dressing, made up of: 300 mg dm⁻³ N, 200 mg dm⁻³ P, 300 mg dm⁻³ K, 50 mg dm⁻³ S, 0.5 mg dm⁻³ B, 1.5 mg dm⁻³ Cu, 3.0 mg dm⁻³ Mn, 0.1 mg dm⁻³ Mo, 5.0 mg dm⁻³ Zn and 5.0 mg dm⁻³ Fe; added as solutions of NH₄H₂PO₄, K₂SO₄, KNO₃, NH₄NO₃, H₃BO₃, CuSO₄·5H₂O, ZnSO₄·7H₂O, FeSO₄·7H₂O, MnCl₂·4H₂O and H₂MoO₄·H₂O (pa grade). After each cut, fertilization as top dressing was performed, made up of 240, 170 and 60 mg dm⁻³ of N, K and S, respectively, added as solutions of NH₄H₂PO₄, K₂SO₄ and KNO₃ (pa grade). During the experimental period, the soil water content was maintained at around 60 % of the total pore volume by periodically weighing the pots and adding deionized water to compensate for weight loss.

Three plants per pot of 'Vitoria' palisade grass [*Urochloa brizantha* (C. Hochstetter ex A. Rich.) R. Webster] were cultivated in 4 dm³ of soil, which were cut during the pre-flowering developmental phases (90 and 140 days after the seedling emergence). After each cutting, shoots were dried for 72 h at 60-65 °C in a forced draught oven, weighed (to obtain dry matter) and triturated in a Wiley-type mill. Biomass obtained from the two cuts were combined to determine the concentration of crude protein (CP) (Silva, 1998), Si (Gallo and Furlani, 1978) and nutrients (Malavolta et al., 1997). Accumulation of Si and nutrients in plant shoots was measured, from the relationship between biomass production and concentrations of Si and nutrients, and quantification of the neutral detergent fiber (NDF) and acid detergent fiber (ADF) were carried out according to van Soest (1994). From these values, the dry matter ingestion (DMI = 120 / %NDF), the dry matter digestibility [DMD = 88.9 (0.779 × %ADF)] and the relative feed value (RFV = DMD × DMI / 1.29) of the forage were estimated according to *American Forage and Grassland Council* (Ensminger, 1993).

Data were submitted to the variance analysis by the F test ($p \leq 0.05$) using the SISVAR software (Ferreira, 2008). When significant, the effect of the calcium silicate on the variables was verified by polynomial regression analysis.

Results and Discussion

Dry matter production was similar for the first and second cuts and the total of the two cuts (TDM) had similar behavior. In general, the 50 % substitution of CaCO₃ by CaSiO₃ promoted higher benefits in the production of TDM, which was increased by 2.6, 15.2 and 19.5 % for the LVAd-1, LVAd-2 and LVd soils, respectively, compared to the treatments that only received CaCO₃ (Figure 1). Most likely the increase in biomass, particularly for the plants cultivated on LVAd-2 and LVd, was due to the silicon (Si) added to the soil by CaSiO₃. Marschner (1995) found Si does not have a defined metabolic role in the plants, however, its action provokes indirect effects, which, when combined can contribute to a higher vegetative production. Silicon accumulation in epidermal cells of the plant shoot promotes a better leaf opening angle, making them more erect, reduces self-shading favoring a better light use and it elevates the photosynthetic rate, culminating in higher biomass production




Figure 1 – Production of total dry matter (TDM) for palisade grass shoots cultivated on LVAd-1, LVAd-2 and LVd soils, as a function of substitution proportions of the calcium carbonate by calcium silicate. * and ** = $p \leq 0.05$ and $p \leq 0.01$, respectively.

(Yoshida et al., 1962). Other beneficial effects of Si in grasses have also been noted, such as reduced transpiration (Dayanandam et al., 1983) and nutrient imbalance (Marschner, 1995; McKeague and Cline, 1963), and increased protection against UV radiation (Tisdale et al., 1993), pest and disease resistance (Rodrigues et al., 2004) and antioxidant enzyme activity (Liang et al., 2003).

In general, the effect of treatments on biomass production was less evident for LVAd-1, probably due its lower clay content (Table 1). Dry matter production by the control (0 % CaCO₃ and 0 % CaSiO₃) was inferior to the other treatments, which shows the importance of soil correction for palisade grass production.

The concentrations and accumulations of calcium (Ca) and magnesium (Mg) in palisade grass shoots (Figure 2 and Table 3) increased considerably with increasing proportions of CaSiO₃ up to an average of 65 %. This behavior can be due to the higher reactivity of the silicates (Lackner, 2002; Ramos et al., 2006). Alcarde and Rodella (2003) found CaSiO₃ to be 6.78 times more soluble than CaCO₃. These factors could have contributed to increasing the absorption of these nutrients due to higher availability of Ca and Mg to the palisade grass. Kaya et al. (2006) and Mali and Aery (2008) verified increases in the concentrations of Ca in corn (*Zea mays*) and cowpea (*Vigna unguiculata*), respectively, with application of silicates to the soil, in agreement with the results obtained in this work.

The accumulations of Ca and Mg in palisade grass shoots, in the soil without correction (0 % CaCO₃ and 0 % CaSiO₃), was generally similar to treatments that received only CaCO₃. This shows that soil correction with only CaCO₃ had little influence on the Ca and Mg absorption by the palisade grass. Concentrations of exchangeable Ca²⁺ and Mg²⁺ in the three soils without correction (Table 1) were in the ranges considered by van Raij et al. (1996) as average




Figure 2 – Concentration and accumulation of macronutrients for palisade grass shoots cultivated on LVAd-1, LVAd-2 and LVd soils, as a function of substitute proportions of the calcium carbonate by calcium silicate.

(4-7 mmol_c of Ca⁺² dm⁻³) and low (0-4 mmol_c of Mg⁺² dm⁻³), respectively.

The concentration and the accumulation of nitrogen (N), potassium (K) and sulfur (S) in the shoot of the palisade grass (Figure 2 and Table 3), when cultivated in LVd, decreased as the CaSiO₃ proportions increased, until an average value of 70 % of CaSiO₃, when it tended to stabilize. Calcium silicate at 70 % of the correction for LVd reduced the concentrations of N, K and S to 58, 49 and 25 %, respectively, and their accumulations to 58, 49 and 9 %, respectively. However, for the plants cultivated in LVAd-1 and LVAd-2, the concentrations and accumulations of these nutrients were mostly unaffected. Conversely, S increased in shoots as CaSiO₃ increased. LVd has a higher cation exchange capacity (CEC) and levels of organic matter, gibbsite, iron oxides and clay, compared to the other appraised soils (Table 1). This partly explains

the differentiated behavior of the plants as to the N, K and S absorption as a function of soil type. For instance, the higher CEC of LVd demanded a higher amount of correctives to elevate the base saturation to 60 %. This resulted in a larger amount of Si being added with CaSiO₃. Wallace (1989) reports that for the absorption sites of the plant there is competition among the anions H₃SiO₄⁻ and NO₃⁻. The lower accumulation of K and S observed with increased CaSiO₃ proportions can likely be explained by the higher solubility of CaSiO₃ compared to CaCO₃, which increases soil solution Ca²⁺ and Mg²⁺, antagonism for the absorption of K⁺ (Malavolta et al., 1997), and low solubility SO₄²⁻. To reiterate, accumulations of N and K in the shoot of the palisade grass were lower when soils did not receive correctives, showing the influence of correctives on N and K absorption by the palisade grass, regardless of soil type.

Table 3 – Regression equations for concentration and accumulation of macronutrients in palisade grass shoots cultivated on LVAd-1, LVAd-2 and LVd soils, as a function of substitute proportions of the calcium carbonate by calcium silicate.

Factor	Soils	Regression equation for concentration		Regression equation for accumulation	
N	LVAd-1	$\hat{y} = 13.3848 - 0.0780x + 0.000884x^2$	$R^2 = 87.28^{**}$	$\hat{y} = 2.0024 - 0.0092x + 0.000102x^2$	$R^2 = 75.52^*$
N	LVAd-2	$\hat{y} = 14.3781 - 0.0793x + 0.000617x^2$	$R^2 = 78.34^*$	$\hat{y} = 1.9117 - 0.00134x$	$R^2 = 50.62_{ns}$
N	LVd	$\hat{y} = 25.0981 - 0.4196x + 0.003017x^2$	$R^2 = 86.38^{**}$	$\hat{y} = 2.9056 - 0.0374x + 0.000244x^2$	$R^2 = 83.46^{**}$
P	LVAd-1	$\hat{y} = 2.4397 - 0.0080x + 0.000086x^2$	$R^2 = 83.51_{ns}$	$\hat{y} = 0.3649 - 0.0007x + 0.000007x^2$	$R^2 = 47.27_{ns}$
P	LVAd-2	$\hat{y} = 2.1114 - 0.0033x + 0.000037x^2$	$R^2 = 28.64_{ns}$	$\hat{y} = 0.2815 + 0.0011x - 0.000009x^2$	$R^2 = 50.02_{ns}$
P	LVd	$\hat{y} = 2.5477 - 0.0243x + 0.000189x^2$	$R^2 = 94.40^{**}$	$\hat{y} = 0.2948 - 0.0012x + 0.000007x^2$	$R^2 = 85.74_{ns}$
K	LVAd-1	$\hat{y} = 8.0114 - 0.0497x + 0.000297x^2$	$R^2 = 82.13_{ns}$	$\hat{y} = 1.1976 - 0.0061x + 0.000028x^2$	$R^2 = 80.06_{ns}$
K	LVAd-2	$\hat{y} = 2.4397 - 0.0080x + 0.000086x^2$	$R^2 = 91.72^*$	$\hat{y} = 1.3389 - 0.0110x + 0.000067x^2$	$R^2 = 79.72^*$
K	LVd	$\hat{y} = 16.0114 - 0.1897x + 0.001097x^2$	$R^2 = 93.86^{**}$	$\hat{y} = 1.8058 - 0.009041x$	$R^2 = 91.34^{**}$
Ca	LVAd-1	$\hat{y} = 2.1517 + 0.0807x - 0.000519x^2$	$R^2 = 91.44^{**}$	$\hat{y} = 0.3230 + 0.0130x - 0.000089x^2$	$R^2 = 90.77^{**}$
Ca	LVAd-2	$\hat{y} = 1.2679 + 0.1817x - 0.001461x^2$	$R^2 = 92.54^{**}$	$\hat{y} = 0.1696 + 0.0288x - 0.000237x^2$	$R^2 = 94.23^{**}$
Ca	LVd	$\hat{y} = 0.71196 + 0.1319x - 0.000748x^2$	$R^2 = 97.10^{**}$	$\hat{y} = 0.0783 + 0.0197x - 0.000126x^2$	$R^2 = 97.08^{**}$
Mg	LVAd-1	$\hat{y} = 0.5194 + 0.0077x - 0.000056x^2$	$R^2 = 75.55^{**}$	$\hat{y} = 0.0777 + 0.0013x - 0.000010x^2$	$R^2 = 74.45^{**}$
Mg	LVAd-2	$\hat{y} = 0.3712 + 0.0239x - 0.000195x^2$	$R^2 = 86.44^{**}$	$\hat{y} = 0.0502 + 0.0039x - 0.000033x^2$	$R^2 = 88.53^{**}$
Mg	LVd	$\hat{y} = 0.2619 + 0.0107x - 0.000078x^2$	$R^2 = 91.95^{**}$	$\hat{y} = 0.0304 + 0.0017x - 0.000014x^2$	$R^2 = 92.93^{**}$
S	LVAd-1	$\hat{y} = 0.8760 + 0.003073x$	$R^2 = 82.09^{**}$	$\hat{y} = 0.1344 + 0.000411x$	$R^2 = 74.88^{**}$
S	LVAd-2	$\hat{y} = 0.7782 + 0.0107x - 0.000123x^2$	$R^2 = 90.43^{**}$	$\hat{y} = 0.1042 + 0.0032x - 0.000025x^2$	$R^2 = 92.51^{**}$
S	LVd	$\hat{y} = 1.4918 - 0.0117x + 0.000091x^2$	$R^2 = 72.31^{**}$	$\hat{y} = 0.1709 - 0.000239x$	$R^2 = 55.68^*$

* and ** = $p \leq 0.05$ and $p \leq 0.01$, respectively. ns = Not significant ($p > 0.05$).

As for the phosphorus (P), CaSiO_3 proportions, up to an average of 64 % CaSiO_3 , decreased P concentration in palisade grass shoots for the LVd (Figure 2 and Table 3). However, the P accumulation (Figure 2 and Table 3) was not significant ($p > 0.05$) among treatments, showing that a lower P concentration was likely due to the “dilution” effect, which is confirmed by the higher shoot dry matter production observed for the 60 % CaSiO_3 treatment (Figure 1). For example, when plant production increases, the “dilution” effect for some nutrients can occur (Crusciol et al., 2008; Marschner, 1995; Miyauchi et al., 2008), which is the reduction of their concentration in the tissue without an alteration in the absorbed quantity. Still, the concentration of P in the plants cultivated on LVAd-1 and LVAd-2 were not different either among treatments. This shows that for P, the substitution of CaCO_3 by CaSiO_3 , in any proportion, does not influence its absorption by palisade grass. Tokura et al. (2007) also verified no effect of silicon doses on P uptake for upland rice (*Oryza sativa* L.) grown on dystroferic Rhodic Acrustox and orthic Ustic Quartzipsamments.

The accumulation of P was much lower in relation to the other plants for palisade grass cultivated in soil without correction. The elevation of the pH promoted by soil correction increases the concentration and activity of the OH^- ion in solution, generating negative charges for the deprotonation of hydroxyls in the exposed clay and organic matter. Thus, the repulsion between orthophosphate ions and the soil mineral particles increases, allowing more P to stay in solution and remain available for plant uptake, especially in soils with high oxide and clay levels. The precipita-

tion of Fe and Al reduces the formation of low solubility P-Fe and P-Al compounds (Novais and Smyth, 1999).

Concentrations and accumulations of the cationic micronutrients in the shoot of the palisade grass (Figure 3 and Table 4) tended to reduce with the elevation of the CaSiO_3 proportions, particularly for Zn. Besides the direct effect of silicate, the increased absorption of Ca^{2+} and Mg^{2+} by palisade grass with increased proportions of CaSiO_3 could have been one of the causes of lower Zn^{2+} absorption, as Malavolta et al. (1997) found competitive inhibition among those nutrients for plant absorption sites. Concentration and accumulation of Cu were decreased only for plants cultivated on LVd. Malavolta et al. (1997) also mentioned Ca^{2+} can impede the exaggerated absorption of Cu^{2+} . The effects of the treatments on the concentrations and accumulations of Mn and Fe were also influenced by the soil type. Decreased accumulations of these nutrients in the shoot of the plants were significant only for those cultivated on LVAd-1. Silicon can increase the parenchyma quantity and diameter in the grasses elevating the oxidant power of the roots, therefore reducing the availability of Mn^{2+} and Fe^{2+} (Korndörfer et al., 1999; Marschner, 1995).

Boron concentration and accumulation (Figure 3 and Table 4) increased as CaSiO_3 proportions were elevated in the three soils, until an average value of 57 % CaSiO_3 , thereafter decreasing in plants cultivated on LVd and LVAd-1 soils. In comparison, the CaSiO_3 proportion of 57 % in the correction of LVAd-1, LVAd-2 and LVd promoted increases of 71, 30 and 53 % in the B concentration, respectively, and 74, 34 and 85 % in the accumulation, respectively, in shoots. Marschner (1995) mentioned H_4SiO_4 and H_3BO_3 , which are




Figure 3 – Concentration and accumulation of micronutrients and Si for palisade grass shoots cultivated on LVAd-1, LVAd-2 and LVd soils, as a function of substitute proportions of the calcium carbonate by calcium silicate.

the main forms of Si and B absorbed by plants, present some similarities as to their behavior in soil solution and plant tissue; both acids are very weak in aqueous solutions and, in the plant, they concentrate in the cell wall, interacting with pectins and polyphenols.

Silicon concentration and accumulation (Figure 3 and Table 4) increased linearly with the elevation of the CaSiO_3 proportions in the three soils. Plants cultivated on LVd and LVAd-2 had a higher rate of increase in Si content with increased CaSiO_3 , compared to those cultivated on LVAd-1. The range of Si concentration was between 5 to 23 g kg^{-1} of Si, which is equal to 1.1-4.9 % of SiO_2 , a range considered appropriate by Marschner (1995) for grasses grown on non-flooded soil.

Concentration and accumulation of Si in the soil without correction were similar to those of plants that received only CaCO_3 (0 % CaSiO_3 treatment), regardless of the soil type. As a result, it is explicit that the additional Si absorbed

by palisade grass in treatments that received CaSiO_3 originated from this source. These results corroborate those of Melo et al. (2003) who also verified increases in concentration and accumulation of Si in the palisade grass shoots with the application of this element to the soil.

The substitution of the CaCO_3 by CaSiO_3 promoted decreased crude protein (CP) concentration in palisade grass shoots for plants cultivated on the LVd. The highest CaSiO_3 proportion also provided reduced NDF and ADF, consequently, resulting in an increase of the estimated values of dry matter digestibility (DMD), dry matter ingestion (DMI) and relative feed value (RFV) of the forage (Figure 4). According to Ensminger (1993) DMD and DMI values of forage are correlated with the concentrations of ADF and NDF, respectively.

Thus, the increased concentration and accumulation of Si (Figures 2 and 3) did not result in reduction of forage feed quality. To the contrary, an increase in RFV was observed

Table 4 – Regression equations for concentration and accumulation of micronutrients and Si in palisade grass shoots cultivated on LVAd-1, LVAd-2 and LVd soils, as a function of substitute proportions of the calcium carbonate by calcium silicate.

Factor	Soils	Regression equation for concentration	Regression equation for accumulation
Si	LVAd-1	$\hat{y} = 5.7469 + 0.034235x$ $R^2 = 88.17^{**}$	$\hat{y} = 0.8853 + 0.0047x$ $R^2 = 83.00^*$
Si	LVAd-2	$\hat{y} = 5.3067 + 0.126664x$ $R^2 = 87.64^{**}$	$\hat{y} = 0.7781 + 0.0185x$ $R^2 = 88.99^{**}$
Si	LVd	$\hat{y} = 8.7785 + 0.136692x$ $R^2 = 94.80^{**}$	$\hat{y} = 1.1360 + 0.0176x$ $R^2 = 93.17^{**}$
B	LVAd-1	$\hat{y} = 3.2883 + 0.0944x - 0.000937x^2$ $R^2 = 98.62^{**}$	$\hat{y} = 0.4935 + 0.0150x - 0.000151x^2$ $R^2 = 97.25^{**}$
B	LVAd-2	$\hat{y} = 7.32427 + 0.039041x$ $R^2 = 82.53^{**}$	$\hat{y} = 0.0388 + 0.0062x$ $R^2 = 95.30^{**}$
B	LVd	$\hat{y} = 8.6814 + 0.1406x - 0.001059x^2$ $R^2 = 79.38^{**}$	$\hat{y} = 0.9917 + 0.0279x - 0.000231x^2$ $R^2 = 96.79^{**}$
Cu	LVAd-1	$\hat{y} = 4.2037 - 0.0185x + 0.000157x^2$ $R^2 = 56.02^{ns}$	$\hat{y} = 0.6431 - 0.0020x + 0.000014x^2$ $R^2 = 61.01^{ns}$
Cu	LVAd-2	$\hat{y} = 3.5069 + 0.0041x - 0.000062x^2$ $R^2 = 23.82^{ns}$	$\hat{y} = 0.4691 + 0.0032x - 0.000032x^2$ $R^2 = 51.67^{ns}$
Cu	LVd	$\hat{y} = 6.5403 - 0.0723x + 0.000571x^2$ $R^2 = 98.18^{**}$	$\hat{y} = 0.7600 - 0.0047x + 0.000031x^2$ $R^2 = 97.92^*$
Fe	LVAd-1	$\hat{y} = 71.5174 + 0.2407x - 0.004383x^2$ $R^2 = 90.02^{ns}$	$\hat{y} = 10.7065 + 0.0506x - 0.000826x^2$ $R^2 = 91.28^*$
Fe	LVAd-2	$\hat{y} = 84.2069 - 0.5941x + 0.005090x^2$ $R^2 = 97.90^{ns}$	$\hat{y} = 11.2636 - 0.0295x + 0.000236x^2$ $R^2 = 57.33^{ns}$
Fe	LVd	$\hat{y} = 95.3400 - 1.0723x + 0.008232x^2$ $R^2 = 72.80^*$	$\hat{y} = 11.0183 - 0.0692x + 0.000444x^2$ $R^2 = 62.83^{ns}$
Mn	LVAd-1	$\hat{y} = 278.0420 - 3.4624x + 0.030440x^2$ $R^2 = 93.71^{**}$	$\hat{y} = 41.6557 - 0.4835x + 0.004105x^2$ $R^2 = 91.49^{**}$
Mn	LVAd-2	$\hat{y} = 97.8774 - 1.4099x + 0.010945x^2$ $R^2 = 80.44^{ns}$	$\hat{y} = 13.0046 - 0.1427x + 0.001061x^2$ $R^2 = 74.42^{ns}$
Mn	LVd	$\hat{y} = 158.7883 - 2.2849x + 0.020091x^2$ $R^2 = 71.75^*$	$\hat{y} = 18.1686 - 0.1720x + 0.001496x^2$ $R^2 = 51.46^{ns}$
Zn	LVAd-1	$\hat{y} = 17.2540 - 0.065800x$ $R^2 = 96.90^{**}$	$\hat{y} = 2.6259 - 0.010392x$ $R^2 = 98.02^{**}$
Zn	LVAd-2	$\hat{y} = 11.0620 - 0.04400x$ $R^2 = 99.62^{**}$	$\hat{y} = 1.5587 - 0.005481x$ $R^2 = 84.88^{**}$
Zn	LVd	$\hat{y} = 11.1003 - 0.1841x + 0.001347x^2$ $R^2 = 84.00^{**}$	$\hat{y} = 1.2780 - 0.01618x + 0.000108x^2$ $R^2 = 81.09^*$

* and ** = $p \leq 0.05$ and $p \leq 0.01$, respectively. ns = Not significant ($p > 0.05$).

with up to 50 % of substitution of the CaCO_3 by CaSiO_3 . These results are corroborated by those of Minson (1990) and van Soest (1994) who did not find negative correlation between Si concentration and forage digestibility. Forage quality is directly related to its nutritional composition, anti-nutritional factors and those which are usually involved in plant protection against the predation and degradation (van Soest, 1994). Garleb et al. (1988) found decreased forage digestibility as the lignin concentration increased.

There is no consensus regarding the effect of Si on forage quality (van Soest, 1994), due to the fact that plants with different Si concentrations also have different composition of other components such as lignin concentration, carbon to nitrogen ratio and differentiated arrangements of cellulose fibrils, which could affect the forage digestibility. Minson (1971) found that Si caused lignin reduction in leaf tissues due to the similar role of both in strengthening cell walls.

The presence of Si in the forage is of great interest because it is considered an essential element for animals being a component of mucopolysaccharides in connective tissues. In herbivores the ingestion of high amounts of phytoliths can cause excessive abrasion in the rumen wall and the dissolved Si can form kidney concretions, causing serious damage to the livestock (Marschner, 1995).

References

Alcarde, J.C.; Rodella, A.A. 1996. The calcium carbonate equivalent of agricultural liming materials. *Scientia Agricola* 53: 6-12. (in Portuguese, with abstract in English).

- Alcarde, J.C.; Rodella, A.A. 2003. Quality and legislation of fertilizers and correctives. p. 291-334. In: Curi, N.; Marques, J.J.; Guilherme, L.R.G.; Lima, J.M.; Lopes, A.S.; Alvares Venegas, V.H, eds. Topics in soil science. Sociedade Brasileira de Ciência do Solo, Viçosa, MG, Brazil. (in Portuguese).
- Alvarez V, V.H.; Novais, R.F.; Dias, L.E.; Oliveira, J.A. 2000. Determination and utilization of solution equilibrium phosphorus. *Boletim Informativo da Sociedade Brasileira de Ciência do Solo* 25: 27-32. (in Portuguese).
- Ávila, F.W.; Baliza, D.P.; Faquin, V.; Araújo, J.L.; Ramos, S.J. 2010. Silicon-nitrogen interaction in rice cultivated under nutrient solution. *Revista Ciência Agronômica* 41: 184-190. (in Portuguese, with abstract in English).
- Crusciol, C.A.C.; Arf, O.; Soratto, R.P.; Mateus, G.P. 2008. Grain quality of upland rice cultivars in response to cropping systems in the Brazilian Tropical Savanna. *Scientia Agricola* 65: 468-473.
- Dayanandam, P.; Kaufman, P.B.; Frankin, C.I. 1983. Detection of silica in plants. *American Journal of Botany* 70: 1079-1084.
- Empresa Brasileira de Pesquisa Agropecuária [EMBRAPA]. 1997. *Methods of Soil Analysis*. 2ed. Centro Nacional de Pesquisa de Solos, Rio de Janeiro, RJ, Brazil. (in Portuguese).
- Empresa Brasileira de Pesquisa Agropecuária [EMBRAPA]. 2006. *Brazilian System of Soil Classification*. 2ed. Centro Nacional de Pesquisa de Solos, Rio de Janeiro, RJ, Brazil. (in Portuguese).
- Ensminger, M.E. 1993. *Dairy Cattle Science*. 3ed. Interstate, Danville, IL, USA.
- Ferreira, D.F. 2008. SISVAR: a program for statistical analysis and teaching. *Revista Symposium* 6: 36-41. (in Portuguese, with abstract in English).
- Gallo, J.R.; Furlani, P.R. 1978. Determination of silicon in plants by the molybdenum-blue colorometric method. *Bragantia* 37: 5-11. (in Portuguese, with abstract in English).
- Garleb, K.A.; Fahey Jr., G.N.; Lewis, S.M.; Kerley, S.M.; Montgomery, L. 1988. Chemical composition and digestibility of fiber fractions of certain by-product feedstuffs fed to ruminants. *Journal of Animal Science* 66: 2650-2662.




Figure 4 – Concentration of crude protein (CP), neutral detergent fiber (NDF) and acid detergent fiber (ADF), and dry matter digestibility (DMD), dry matter ingestion (DMI, in % of body weight) and relative feed value (RFV) for palisade grass shoots, as a function of substitute proportions of the calcium carbonate by calcium silicate. ** and ns = $p \leq 0.01$ and not significant ($p > 0.05$), respectively.

Jones, L.H.P.; Handreck, K.A. 1967. Silica in soils, plant and animals. *Advances in Agronomy* 19: 107-149.

Kaya, C.; Tuna, L.; Higgs, D. 2006. Effect of silicon on plant growth and mineral nutrition of maize grown under water-stress conditions. *Journal of Plant Nutrition* 29: 1469-1480.

Korndörfer, G.H.; Arantes, V.A.; Corrêa, G.F.; Snyder, G.H. 1999. Effect of calcium silicate on soil silicon content and upland rice grain yield. *Revista Brasileira de Ciência do Solo* 23: 635-641. (in Portuguese, with abstract in English).

Klug, H.P.; Alexander, L.E. 1974. *X-ray Diffraction Procedures for Polycrystalline and Amorphous Materials*. John Wiley, New York, NY, USA.

Lackner, K.S. 2002. Carbonate chemistry for sequestering fossil carbon. *Annual Review of Energy and the Environment* 27: 193-232.

Liang, Y.C.; Chen, Q.; Liu, Q.; Zhang, W.H.; Ding, R.X. 2003. Exogenous silicon (Si) increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). *Journal of Plant Physiology* 160: 1157-1164.

Malavolta, E.; Vitti, G.C.; Oliveira, S.A. 1997. *Assessment of Nutritional Status of Plants: Principles and Applications*. 2ed. Potafos, Piracicaba, SP, Brazil. (in Portuguese).

Mali, M.; Aery, N.C. 2008. Silicon effects on nodule growth, dry-matter production, and mineral nutrition of cowpea (*Vigna unguiculata*). *Journal of Plant Nutrition and Soil Science* 171: 835-840.

Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. Academic Press, London, England.

Matoh, T.; Kairusmee, P.; Tokahashi, E. 1986. Salt-induced damage to rice plants and alternation effect of silicate. *Soil Science and Plant Nutrition* 32: 295-304.

McKeague, J.A.; Cline, M.G. 1963. Silica in the soil. *Advances in Agronomy* 15: 339-396.

Melo, S.P.; Korndörfer, G.H.; Korndörfer, C.M.; Lana, R.M.Q.; Santana, D.G. 2003. Silicon accumulation and water deficit tolerance in *Brachiaria* grasses. *Scientia Agricola* 60: 755-759.

Mengel, K.; Kirkby, E.A. 2001. *Principles of Plant Nutrition*. 5ed. Kluwer Academic, Norwell, MA, USA.

Mehra, O.P.; Jackson, M.L. 1960. Iron oxide removal from soils and clays by a dithionite-citrate system buffered with sodium bicarbonate. *Clays and Clay Minerals* 7: 317-327.

Minson, D.J. 1971. Influence of lignin and silicon on a summative system for assessing the organic matter digestibility of Panicum. *Australian Journal of Agricultural Research* 22: 589-598.

- Minson, J.D. 1990. Forages in Ruminant Nutrition. Academic Press, New York, NY, USA.
- Miyachi, M.Y.H.; Lima, D.S.; Nogueira, M.A.; Lovato, G.M.; Murate, L.S.; Cruz, M.F.; Ferreira, J.M.; Zangaro, W.; Andrade, G. 2008. Interactions between diazotrophic bacteria and mycorrhizal fungus in maize genotypes. *Scientia Agricola* 65: 525-531.
- Novais, R.F.; Smyth, T.J. 1999. Phosphorus in Soil and Plants under Tropical Conditions. Universidade Federal de Viçosa, Viçosa, MG, Brazil. (in Portuguese).
- Ramos, L.A.; Nolla, A.; Korndörfer, G.H.; Pereira, H.S.; Camargo, M.S. de. 2006. Reactivity of soil acidity correctives and conditioners in lysimeters. *Revista Brasileira de Ciência do Solo* 30: 849-857. (in Portuguese, with abstract in English).
- Reis, G.L.; Lana, A.M.Q.; Maurício, R.M.; Lana, R.M.Q.; Machado, R.M.; Borges, I.; Quinzeiro Neto, T. 2010. Influence of trees on soil nutrient pools in a silvopastoral system in the Brazilian Savannah. *Plant and Soil* 329: 185-193.
- Rodrigues, F.A.; McNally, D.J.; Datnoff, L.E.; Jones, J.B.; Labbe, C.; Benhamou, N.; Menzies, J.G. 2004. Silicon enhances the accumulation of dipentenoid phytoalexins in rice: a potential mechanism for blast resistance. *Phytopathology* 94: 177-183.
- Schwertmann, U. 1964. The differentiation of iron oxides in soils by extraction with ammonium oxalate solution. *Zeitschrift Pflanzenernähr Düng Bodenkd* 105: 194-202. (in German).
- Silva, D.J. 1998. Food Analysis: Chemical and Biological Methods. Universidade Federal de Viçosa, Viçosa, MG, Brazil. (in Portuguese).
- Tisdale, S.L.; Nelson, W.L.; Beaton, J.D.; Havlin, J.L. 1993. Soil Fertility and Fertilizers. 5ed. Macmillan, New York, NY, USA.
- Tokura, A.M.; Furtini Neto, A.E.; Curi, N.; Carneiro, L.F.; Alovisei, A.A. 2007. Silicon and phosphorus in soils cultivated with upland rice. *Acta Scientiarum Agronomy* 29: 9-16. (in Portuguese, with abstract in English).
- Van Raij, B.; Cantarella, H.; Quaggio, J.A.; Ferreira, H.E.; Lopes, A.S.; Bataglia, O.C. 1987. Soil Chemical Analysis for Fertility Purposes. Fundação Cargill, Campinas, SP, Brazil. (in Portuguese).
- Van Raij, B.; Cantarella, H.; Quaggio, J.A.; Furlani, A.M.C. 1996. Fertilization and Liming Recommendation for the State of São Paulo. 2ed. Instituto Agrônomo de Campinas, Campinas, SP, Brazil. (in Portuguese).
- Van Soest, P.J. 1994. Nutritional Ecology of the Ruminant. 2ed. Cornell University Press, Ithaca, NY, USA.
- Wallace, A. 1989. Relationships among nitrogen, silicon, and heavy metal uptake by plants. *Soil Science* 147: 457-460.
- Yoshida, S.; Ohnishi, Y.; Kitagishi, K. 1962. Chemical forms, mobility and deposition of silicon in rice plant. *Soil Science and Plant Nutrition* 8: 15-21.

Received April 26, 2010

Accepted January 20, 2011